
Automatic classification of birds calls and songs from Provence

Olivier Dufour*^{†1}, Hervé Glotin¹, Thierry Artières², Yves Bas³, Pascale Giraudet⁴, and Vincent Delcourt³

¹Laboratoire des Sciences de l'Information et des Systèmes (LSIS) – UMR 7296 – Université de Toulon
83957 La Garde, France

²Laboratoire d'Informatique de Paris 6 (LIP6) – Université Pierre et Marie Curie (UPMC) - Paris VI,
CNRS : UMR7606 – 4 Place JUSSIEU 75252 PARIS CEDEX 05, France

³Biotope – RD – 22 Bd Maréchal Foch BP 58 34140 MEZE, France

⁴Université Sud Toulon Var (USTV) – UFR Sc et Tech – Avenue de l'Université - BP 20132 - 83957 La
Garde cedex, France

Abstract

Neural Information Processing Scaled for Bioacoustics bird challenge consisted in automatically identify 87 classes present in thousand audio recordings, collected in many places of Provence with Song Meter recorders. The difficulty of this task lies in the large number of classes, various background noises and simultaneously vocalising animals. Among 32 international teams, best Area Under the Curve score of the official benchmark has been 91.7%, which corresponds to a mean recall of 50% (all classes confunded).

Our objective is to increase performances to get an operational system. It implies to reach a 80 % mean recall.

For this, we decided to merge winning models of NIPS4B, test new improvements and add training databases : xeno-canto.org (161 605 recordings) and Animal Sound Archive (Natural History Museum of Berlin, 5 209 recordings).

We succeeded in increasing mean recall and precision. Plus, our recall and precision calculations are statistically solid: for each class, the number of test files is important and equivalent.

In february 2014, we began a twelve monthes continuous acoustic recording experience in 10 different sites: natural habitats (wetlands, oak groves, ...) and semi natural habitats (melange between garrigue and grapevines) affected by anthropogenic activities whose some of them are suspected to be threats (windfirms) for avifauna. In conclusion, our algorithms are now enough good to:

- precisely measure temporal evolution of acoustic activity of 50 species of birds from Eastern Palaeartic. As far as we know, unbiased studies of daily and annual phenology of migratory and non migratory bird species are completely unseen. Plus, as we discriminate song and

*Speaker

[†]Corresponding author: olivierlouis.dufour@gmail.com

call for most of birds, we can monitor the calls/songs ratio during bird migration and the rest of the year.

- check efficacy of bird acoustic repellent systems for windfirms.

Keywords: automatic, birds, songs, classification